



RankenJordan
PEDIATRIC BRIDGE HOSPITAL

Intermittent Clean Catheterization for Males

What is intermittent cathing?

- Intermittent catheterization (“cathing”) involves a child or caregiver inserting a catheter (a long thin tube) through the urethra (the smooth passage from the bladder to the outside of the body that carries urine) into the bladder to empty urine. Once the bladder is empty the catheter is removed.

Why does my child need intermittent cathing?

- Your child needs intermittent cathing because they cannot empty their bladder fully on their own. This is often caused by damage to the nerves connecting the spinal cord and the sphincter muscles (muscles that tighten to hold urine in or relax to let urine out).
- Emptying the bladder with a catheter throughout the day allows your child to be dry between cathing. This helps prevent urinary tract infections, problems with the kidneys, and other more serious problems.

How often does my child need intermittent cathing?

- Your child needs to be cathed at the following times every day -

- It is important that you cath your child on time and do not skip any scheduled times.
- If you miss any of the above times, cath your child as soon as possible.

How do I perform clean cathing?

1. Gather supplies.
 - a. Catheter—size: _____ French
 - b. Baby wipes or clean cloth with soap and water
 - c. Container for urine
 - d. Water soluble lubricant
2. Move your child into a private place and remove your child's pants and underwear or diaper.
3. Wash your hands.
4. Put on clean gloves.
5. If your child is not circumcised, retract the foreskin.
6. Grasp the shaft of the penis with your non-dominant hand and continue to hold while cathing.
7. Wash the tip of the penis with soap and water or a baby wipe.

Picture A.
8. Pick up the catheter with your dominant hand and lubricate the tip.
9. Place the drainage end of the catheter into the container, such as a urinal, to collect the urine. There is usually a colored, rubber end on the drainage end of the catheter.
10. Locate the urethral opening (hole at the tip of the penis that leads to the bladder). Insert the catheter until urine flows out. Push the catheter in one more inch to make sure it is well in the bladder. *Picture B.*

Picture A.



Picture B.



11. Continue to hold the penis and the catheter in place while the urine flows out.

12. Once the urine stops flowing, gently push on the lower stomach over the bladder to help the bladder to fully empty.

Picture C.

13. When urine stops flowing, pull the catheter out about an inch to see if any more urine flows out. If it does, wait until urine stops flowing before removing the catheter.

14. When there is no more urine coming out of the catheter, pinch or bend the catheter to prevent leakage and backflow of urine into the bladder, and gently pull the catheter out. *Picture D.*

15. Use a wet washcloth or a baby wipe to clean any excess urine off your child's skin.

16. Pour the urine down the toilet. Rinse the supplies you need to keep for later with warm, soapy water. Throw away any other supplies.

Picture C.



Picture D.



When to call the doctor:

- Fever
- Foul-smelling urine
- Blood in the urine
- A lot of sediment (floating particles that eventually settle to the bottom) in the urine (that looks different than usual for your child).
- Pain in the lower back or lower abdomen
- Trouble inserting the catheter
- Pain while catheterizing