Trach 1-2-3 Action



PROBLEM	SIGNS/SYMPTOMS	ACTION 1-2-3		
Plugged Trach Prevention: Humidification Secretion management	 Difficulty breathing Coughing Whistling noise Skin color change Agitation Retractions, nasal flaring 	Suction per care plan/safe suction guide	No improvement? Change trach	No improvement? Call 911
Infection Prevention: Hand hygiene High touch surface cleaning	 Fever Increased/thick secretions Change in secretion color Swelling/redness at stoma Odor 	Document vitals & sick/illness log	Call caregiver & provider	
Bleeding Prevention: Safe suction guide Stoma care	 Light pink/streaky red blood with suctioning Bright red blood = Immediate Action - Call 911 	Monitor closely and document	Notify caregiver & provider	Call 911 for profuse bright red bleeding
Accidental Dislodgement Prevention: Trach tie tightness Tube position Distraction/Redirection	 Tube not in stoma Respiratory distress Skin color change DON'T PANIC	If trach is in: reposition trach, tighten ties and check balloon If trach is out: change trach	If <u>same size</u> doesn't go in, use 1 size smaller trach	If unable to insert <u>any</u> size trach, call 911 and use Ambu bag for rescue breathing

Perform CPR if child ever goes unresponsive with NO PULSE or pulse < 60 (Birth - 12 months) or pulse < 40 (13+ months)

Perform RESCUE breathing if child is NOT BREATHING | Airway Bag should ALWAYS be with child

This document contains information and/or instructional materials developed by Ranken Jordan Pediatric Bridge Hospital. This content does not replace medical advice, diagnosis, or treatment. Talk to your child's health care provider if you have any questions about this document, your child's condition, or your child's treatment plan. If your child is not a Ranken Jordan Pediatric Bridge Hospital patient, this document does not create a doctorpatient relationship between your child and Ranken Jordan Pediatric Bridge Hospital.